

Confidential: Proprietary information of Ivorychem Pte Ltd. Not to be disclosed to third parties without prior consent of Ivorychem Pte Ltd

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.

USE ONLY IN ACCORDANCE WITH INSTRUCTIONS.

KEEP OUT OF REACH OF CHILDREN

ACTIFRON 75 WDG



INGREDIENTS

Halosulfuron.....75%
Other ingredients25%

ACTIFRON 75 WP is a herbicide recommended to control yellow and purple nutsedges and to suppress green kyllinga in bahiagrass, bermudagrass, centipedegrass, St. Augustinegrass, and zoysiagrass, and has as its active ingredient, Halosulfuron.

Trade names for products containing Halosulfuron include Manage.

What is ACTIFRON 75 WDG and how does it work?

Key Benefits of ACTIFRON 75 WDG:

PRECAUTIONS

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse. Keep children or pets away from treated area until dry.

SYMPTOMS OF POISONING

Irritation on skin or eyes.

MEDICAL TREATMENT

No specific antidote is available. Treatment is symptomatic.

FIRST AID

If on skin, remove contaminated clothes. Rinse and then rinse skin immediately with plenty of water and soap for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person for fresh air. If person is not breathing, call for an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, first hold eye open and rinse with plenty of water for 15-20 minutes (remove contact lenses if easily possible). Call poison control center or doctor for treatment advice.

If ingested, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do NOT induce vomiting unless told to do so by poison control center or doctor. Do not give anything to an unconscious person.

DIRECTIONS OF USE

The use rate of ACTIFRON 75 WDG is 0.015 to 0.031 ounces/1000 ft² (= 0.43 to 0.87 grams) or 0.01 to 0.02 ounces ai/1,000 ft.² Apply after nutsedge has 3 to 8 leaves. Use lower rates for light infestations and the higher rate for heavy infestations.

A second application may be needed 6 to 10 weeks after the initial treatment for complete control. Add nonionic surfactant at 0.5 percent with ACTIFRON 75 WP applications.

Grasses should be well established at the time of application. Do not mow turf 2 days before or less than 2 days after ACTIFRON 75 WP applications. Do not apply directly over the top of flowers, ornamentals, shrubs, or other desirable plants. Do not use clippings from treated areas to mulch around vegetables or ornamentals.

Crop	Application rate (pounds per acre treated)
Corn for Forage	0.05
Landscape	0.06
Almonds	0.02
Walnuts	0.03
Corn	0.04
Cantaloupe	0.037
Outdoor Container Nursery	0.05
Uncultivated Agricultural Area	0.06
Pistachios	0.02
Ornamental Turf	0.05
Tomatoes for Processing	0.02
Soil Preplant	0.04
Sorghum	0.04
Cucumbers	0.02
Asparagus	0.47
Rice	0.05

Outdoor Propagation Nursery	0.06
Tomatoes	0.02
Alfalfa for Forage	0.05
Pasture	0.05
Squash	0.02
Melons	0.0077
Watermelon	0.044
Greenhouse Plants	0.41
Pumpkins	0.03
Greenhouse Flowers	0.04

ACTIFRON 75 WP is effective for weed control at 0.018 lb AI/A and grass seed at 0.67 to 1.33 oz/A.

Caution: No more than four applications can be made with the total use rate not exceeding 5.33 oz/A per season.

Amount of formulation per 1,000 sq ft	Amount of formulation per acre	Pounds active ingredient per acre	Precautions and remarks
0.9 g	0.67 to 1.33 oz	0.031 to 0.062	May be applied to bermudagrass, bahiagrass, zoysiagrass, centipedegrass, St. Augustinegrass, creeping bentgrass, Kentucky bluegrass, perennial ryegrass, tall fescue, and fine fescue. Apply broadcast when sedges have reached the three- to eight-leaf stage. Use lower rate for light infestations and higher rate for heavy infestations. A second

			treatment will usually be required 6 to 10 weeks after the initial treatment. Use an 80% active nonionic surfactant at 2 qt per 100 gal of spray solution (0.5% by volume). Do not exceed 1 to 2 pt of surfactant per acre. Do not apply to putting greens. Manage only suppresses green kyllinga.
--	--	--	--

DISPOSAL METHODS

Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE CONDITION

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank.

For More Details including effects on environment email contact@ivorychem.com with Subject "ACTIFRON 75 WDG DETAILS".

More Details:

TOXICOLOGICAL EFFECTS
ECOLOGICAL EFFECTS
ENVIRONMENTAL FATE

PHYSICAL PROPERTIES AND GUIDELINES

PHYSICAL PROPERTIES:

EXPOSURE GUIDELINES:



IVORYCHEM PTE LIMITED
15 Beach Road #02-09
Beach Centre
Singapore 189677
Tel: +65 63377765
Fax: +65 63377730
contact@ivorychem.com
www.ivorychem.com

Company Registration No 200405572W